


Appendix H: Q Distribution Table

How to Use the Q Distribution Table

This table should be used only if the sample sizes in your Tukey's HSD analysis are equal. There are two sections of the table, one for the .05 significance level (H.1) and one for the .01 significance level (H.2). Select a significance level and the corresponding section of the table. Then, find the number of groups (samples) being compared in the top row of the table. This determines the column of the table you will use. Find the df for your data ($df = n - k$, where k is the number of groups [samples] and n is the number of observations in one of the samples) in the left-hand column of the table. This identifies the row of the table you will use. The critical value of Q for the HSD test is found at the intersection of the row and column you have identified. A difference between sample means as large or larger than the HSD you calculate using the table value of Q is significant at the selected level of significance.

Table H.1 Critical Values of Q ($p = 0.05$)

		$(p) = 0.05$								
$df \backslash k$		2	3	4	5	6	7	8	9	10
1		18.0	27.0	32.8	37.1	40.4	43.1	45.4	47.4	49.1
2		6.08	8.33	9.80	10.88	11.73	12.43	13.03	13.54	13.99
3		4.50	5.91	6.82	7.50	8.04	8.48	8.85	9.18	9.46
4		3.93	5.04	5.76	6.29	6.71	7.05	7.35	7.60	7.83
5		3.64	4.60	5.22	5.67	6.03	6.33	6.58	6.80	6.99
6		3.46	4.34	4.90	5.30	5.63	5.90	6.12	6.32	6.49
7		3.34	4.16	4.68	5.06	5.36	5.61	5.82	6.00	6.16
8		3.26	4.04	4.53	4.89	5.17	5.40	5.60	5.77	5.92
9		3.20	3.95	4.41	4.76	5.02	5.24	5.43	5.59	5.74
10		3.15	3.88	4.33	4.65	4.91	5.12	5.30	5.46	5.60
11		3.11	3.82	4.26	4.57	4.82	5.03	5.20	5.35	5.49
12		3.08	3.77	4.20	4.51	4.75	4.95	5.12	5.27	5.39
13		3.06	3.73	4.15	4.45	4.69	4.88	5.05	5.19	5.32
14		3.03	3.70	4.11	4.41	4.64	4.83	4.99	5.13	5.25
15		3.01	3.67	4.08	4.37	4.59	4.78	4.94	5.08	5.20
16		3.00	3.65	4.05	4.33	4.56	4.74	4.90	5.03	5.15
17		2.98	3.63	4.02	4.30	4.52	4.70	4.86	4.99	5.11
18		2.97	3.61	4.00	4.28	4.49	4.67	4.82	4.96	5.07
19		2.96	3.59	3.98	4.25	4.47	4.65	4.79	4.92	5.04
20		2.95	3.58	3.96	4.23	4.45	4.62	4.77	4.90	5.01
24		2.92	3.53	3.90	4.17	4.37	4.54	4.68	4.81	4.92
30		2.89	3.49	3.85	4.10	4.30	4.46	4.60	4.72	4.82
40		2.86	3.44	3.79	4.04	4.23	4.39	4.52	4.63	4.73
60		2.83	3.40	3.74	3.98	4.16	4.31	4.44	4.55	4.65
120		2.80	3.36	3.68	3.92	4.10	4.24	4.36	4.47	4.56
∞		2.77	3.31	3.63	3.86	4.03	4.17	4.29	4.39	4.47

(Continued)

Table H.2 Critical Values of Q ($p = 0.01$)

		$(p) = 0.01$								
$df \backslash k$	2	3	4	5	6	7	8	9	10	
1	90.0	135	164	186	202	216	227	237	246	
2	13.90	19.02	22.56	25.37	27.76	29.86	31.73	33.41	34.93	
3	8.26	10.62	12.17	13.32	14.24	15.00	15.65	16.21	16.71	
4	6.51	8.12	9.17	9.96	10.58	11.10	11.54	11.92	12.26	
5	5.70	6.98	7.80	8.42	8.91	9.32	9.67	9.97	10.24	
6	5.24	6.33	7.03	7.56	7.97	8.32	8.61	8.87	9.10	
7	4.95	5.92	6.54	7.00	7.37	7.68	7.94	8.17	8.37	
8	4.75	5.64	6.20	6.62	6.96	7.24	7.47	7.68	7.86	
9	4.60	5.43	5.96	6.35	6.66	6.91	7.13	7.33	7.49	
10	4.48	5.27	5.77	6.14	6.43	6.67	6.87	7.05	7.21	
11	4.39	5.15	5.62	5.97	6.25	6.48	6.67	6.84	6.99	
12	4.32	5.05	5.50	5.84	6.10	6.32	6.51	6.67	6.81	
13	4.26	4.96	5.40	5.73	5.98	6.19	6.37	6.53	6.67	
14	4.21	4.89	5.32	5.63	5.88	6.08	6.26	6.41	6.54	
15	4.17	4.84	5.25	5.56	5.80	5.99	6.16	6.31	6.44	
16	4.13	4.79	5.19	5.49	5.72	5.92	6.08	6.22	6.35	
17	4.10	4.74	5.14	5.43	5.66	5.85	6.01	6.15	6.27	
18	4.07	4.70	5.09	5.38	5.60	5.79	5.94	6.08	6.20	
19	4.05	4.67	5.05	5.33	5.55	5.73	5.89	6.02	6.14	
20	4.02	4.64	5.02	5.29	5.51	5.69	5.84	5.97	6.09	
24	3.96	4.55	4.91	5.17	5.37	5.54	5.69	5.81	5.92	
30	3.89	4.45	4.80	5.05	5.24	5.40	5.54	5.65	5.76	
40	3.82	4.37	4.70	4.93	5.11	5.26	5.39	5.50	5.60	
60	3.76	4.28	4.59	4.82	4.99	5.13	5.25	5.36	5.45	
120	3.70	4.20	4.50	4.71	4.87	5.01	5.12	5.21	5.30	
∞	3.64	4.12	4.40	4.60	4.76	4.88	4.99	5.08	5.16	

Source: www.stat.wisc.edu/courses/st571-ane/tables/tableQ.pdf.