

Deaths by AIDS in 2006

National Total = 12,113 Deaths*

ALPHA ORDER

| RANK | STATE | DEATHS | % of USA |
|------|----------------|--------|----------|
| 16 | Alabama | 189 | 1.6% |
| 44 | Alaska | 10 | 0.1% |
| 21 | Arizona | 138 | 1.1% |
| 28 | Arkansas | 74 | 0.6% |
| 3 | California | 1,176 | 9.7% |
| 26 | Colorado | 83 | 0.7% |
| 22 | Connecticut | 127 | 1.0% |
| 30 | Delaware | 54 | 0.4% |
| 1 | Florida | 1,753 | 14.5% |
| 5 | Georgia | 684 | 5.6% |
| 40 | Hawaii | 18 | 0.1% |
| 45 | Idaho | 8 | 0.1% |
| 9 | Illinois | 382 | 3.2% |
| 20 | Indiana | 141 | 1.2% |
| 41 | Iowa | 16 | 0.1% |
| 35 | Kansas | 28 | 0.2% |
| 27 | Kentucky | 81 | 0.7% |
| 11 | Louisiana | 338 | 2.8% |
| 42 | Maine | 12 | 0.1% |
| 7 | Maryland | 483 | 4.0% |
| 19 | Massachusetts | 182 | 1.5% |
| 18 | Michigan | 184 | 1.5% |
| 31 | Minnesota | 53 | 0.4% |
| 17 | Mississippi | 187 | 1.5% |
| 23 | Missouri | 120 | 1.0% |
| 47 | Montana | 6 | 0.0% |
| 38 | Nebraska | 24 | 0.2% |
| 28 | Nevada | 74 | 0.6% |
| 42 | New Hampshire | 12 | 0.1% |
| 6 | New Jersey | 539 | 4.4% |
| 34 | New Mexico | 39 | 0.3% |
| 2 | New York | 1,471 | 12.1% |
| 8 | North Carolina | 427 | 3.5% |
| 50 | North Dakota | 2 | 0.0% |
| 15 | Ohio | 232 | 1.9% |
| 25 | Oklahoma | 89 | 0.7% |
| 33 | Oregon | 50 | 0.4% |
| 10 | Pennsylvania | 367 | 3.0% |
| 37 | Rhode Island | 27 | 0.2% |
| 12 | South Carolina | 256 | 2.1% |
| 45 | South Dakota | 8 | 0.1% |
| 12 | Tennessee | 256 | 2.1% |
| 4 | Texas | 1,034 | 8.5% |
| 35 | Utah | 28 | 0.2% |
| 47 | Vermont | 6 | 0.0% |
| 14 | Virginia | 251 | 2.1% |
| 24 | Washington | 100 | 0.8% |
| 39 | West Virginia | 22 | 0.2% |
| 32 | Wisconsin | 52 | 0.4% |
| 49 | Wyoming | 3 | 0.0% |

RANK ORDER

| RANK | STATE | DEATHS | % of USA |
|------|----------------------|--------|----------|
| 1 | Florida | 1,753 | 14.5% |
| 2 | New York | 1,471 | 12.1% |
| 3 | California | 1,176 | 9.7% |
| 4 | Texas | 1,034 | 8.5% |
| 5 | Georgia | 684 | 5.6% |
| 6 | New Jersey | 539 | 4.4% |
| 7 | Maryland | 483 | 4.0% |
| 8 | North Carolina | 427 | 3.5% |
| 9 | Illinois | 382 | 3.2% |
| 10 | Pennsylvania | 367 | 3.0% |
| 11 | Louisiana | 338 | 2.8% |
| 12 | South Carolina | 256 | 2.1% |
| 12 | Tennessee | 256 | 2.1% |
| 14 | Virginia | 251 | 2.1% |
| 15 | Ohio | 232 | 1.9% |
| 16 | Alabama | 189 | 1.6% |
| 17 | Mississippi | 187 | 1.5% |
| 18 | Michigan | 184 | 1.5% |
| 19 | Massachusetts | 182 | 1.5% |
| 20 | Indiana | 141 | 1.2% |
| 21 | Arizona | 138 | 1.1% |
| 22 | Connecticut | 127 | 1.0% |
| 23 | Missouri | 120 | 1.0% |
| 24 | Washington | 100 | 0.8% |
| 25 | Oklahoma | 89 | 0.7% |
| 26 | Colorado | 83 | 0.7% |
| 27 | Kentucky | 81 | 0.7% |
| 28 | Arkansas | 74 | 0.6% |
| 28 | Nevada | 74 | 0.6% |
| 30 | Delaware | 54 | 0.4% |
| 31 | Minnesota | 53 | 0.4% |
| 32 | Wisconsin | 52 | 0.4% |
| 33 | Oregon | 50 | 0.4% |
| 34 | New Mexico | 39 | 0.3% |
| 35 | Kansas | 28 | 0.2% |
| 35 | Utah | 28 | 0.2% |
| 37 | Rhode Island | 27 | 0.2% |
| 38 | Nebraska | 24 | 0.2% |
| 39 | West Virginia | 22 | 0.2% |
| 40 | Hawaii | 18 | 0.1% |
| 41 | Iowa | 16 | 0.1% |
| 42 | Maine | 12 | 0.1% |
| 42 | New Hampshire | 12 | 0.1% |
| 44 | Alaska | 10 | 0.1% |
| 45 | Idaho | 8 | 0.1% |
| 45 | South Dakota | 8 | 0.1% |
| 47 | Montana | 6 | 0.0% |
| 47 | Vermont | 6 | 0.0% |
| 49 | Wyoming | 3 | 0.0% |
| 50 | North Dakota | 2 | 0.0% |
| | District of Columbia | 217 | 1.8% |

Source: U.S. Department of Health and Human Services, National Center for Health Statistics

"National Vital Statistics Reports" (Vol. 57, No. 14, April 17, 2009, <http://www.cdc.gov/nchs/deaths.htm>)

*Final data by state of residence. AIDS is Acquired Immunodeficiency Syndrome. It is a specific group of diseases or conditions which are indicative of severe immunosuppression related to infection with the Human Immunodeficiency Virus (HIV).